

# Length-Weight Relationship of Fishes Caught in the Itaipu Reservoir, Paraná, Brazil

E. Benedito-Cecilio, A.A. Agostinho  
and R.C. Carnelós-Machado Velho

## Abstract

Parameters and related statistics of the length-weight relationship of the form  $W=aL^b$  are presented for 72 species of fish caught in the area of the Itaipu Reservoir in Paraná, Brazil. The  $b$  values varied between 2.34 and 3.35, with the mean  $b=2.986$  (s.d.=0.230) not significantly different from 3.0 (df=7,  $p=0.05$ ).

## Introduction

Basic information such as knowledge on parameters that relate weight to length of fish is scanty, even though it is of great importance in studies on fisheries biology (Vazzoler 1996) and on the evaluation of fish stocks (Entsua-Mensah et al. 1995). Among its most frequent uses, Pauly (1993) notes the importance of length-weight relationships (LWR) in the calculation of the fish's average weight at a certain length class and the conversion of an equation of growth in length into an equation of growth in weight, besides morphological comparisons between populations of the same species or between species. Such applications are evident in the work of Kulbicki et al. (1993), Almeida et al. (1995), Stergiou and Politou (1995), and Petrakis and Stergiou (1995), among others.

The mathematical parameters of the relationship between the length and the weight of fish furnish further information on the weight variation of individuals in

relation to their length (condition factor,  $k$ ). This factor estimates the general well-being of the individual and is frequently used in three cases: (1) comparison of two or more co-specific populations living in similar or different conditions of food, density or climate, among others; (2) determination of the period and duration of gonadal maturation; and (3) observation of the increase or decrease in feeding activity or population changes, possibly due to modifications in food resources (Weatherley and Gill 1987).

Generally, a growth model in fish follows the "cube law" and hence the use of Fulton's condition factor or the isometric factor ( $k = W/L^3$ ), attributing to the weight-length exponent  $b$  a value equal to 3 (Gulland 1983). In this case the body form maintains a constant proportion to length (Weatherley and Gill 1987). However, Braga (1986) states that Fulton's condition factor is only adequate for the comparison of fish of the same size, while the allometric condition factor, which occurs when  $b \neq 3$ , is valid for the

study of fish of any range of length although at the same stage of development. Vazzoler (1996) states that it is inadvisable to use Fulton's condition factor and affirms that the determination of the value of  $b$  is necessary in the use of the allometric condition factor so that results may be reliable.

The present research work aims at making available estimates of the LWR parameters for 72 species of fish caught in the area of the Itaipu Reservoir.

## Materials and Methods

Individual measurements of total length (TL) and standard length (SL) in cm and of total weight (W) in g were obtained from November 1983 to October 1989. Sampling was carried out in 11 stations, 3 in the principal channel of the Itaipu Reservoir (municipalities of Guaíra, Santa Helena and Foz do Iguaçu, state of Paraná, Brazil), 4 in the tributaries of the left shore (Rivers Ocoí, São Francisco Falso, São Francisco Verdadeiro and Arroio Guaçu), 2 in the River

Paraná (one upstream and the other downstream of the Itaipu Reservoir) and 2 in the tributaries of the River Paraná (Rivers Iguatemi and Piquiri) (Fig. 1).

Fishing equipment, exposed for 24 hours and checked at periods of 8-12 hours, consisted of simple gillnets and trammel nets with mesh size from 3 to 16 cm at opposite knots, and boulders with 30 and 50 fishing hooks.

Initially, data analysis consisted of a graphic visualization of the relationship between standard length and total weight. Subse-

quently, regression by the least-squares method for the determination of parameters and related statistics of the LWR whose variables were previously transformed into logarithms ( $1nW = 1n a + b 1n SL$ ). The individual species and average value for  $b$  was tested to verify whether it was significantly different from 3 using t-test at the 0.05 significance level.

### Results and Discussion

The species studied correspond to 96.6% in number and 88.9% in

weight of total fish captured in the area of the Itaipu Reservoir from November 1983 to October 1989 (Benedito-Cecilio 1994). Table 1 gives a summary of the results of the analysis.

The number of individuals varied from 25 in the case of *Pachiuirus bonariensis* to 12 867 for *Auchenipterus nuchalis*. Values of  $r^2$  ranged from 0.81 for *Apareiodon affinis* to 0.99 in the case of *Salminus maxillosus*, *Brycon orbignyianus*, *Serrasalmus spilopleura* and *Loricaria prolixa*, with all regressions highly significant




Fig. 1. Site of sampling stations in the area of the Itaipu Reservoir, Paraná, Brazil.

Table 1. Number of individuals (n), minimum, maximum and mean standard lengths (cm) and parameters of the length-weight relationship for species caught in the Itaipu Reservoir from November 1983 to October 1989. Species with asterisk indicate total length was used rather than standard length; b values in italic.

Family/Species	n	Standard length				Length-weight relationship				
		mean	s.d.	min	max	TL/LS	a	b	s.e.(b)	r <sup>2</sup>
<b>Family Engraulidae</b>										
<i>Lycengraulis grossidens</i>	47	13.1	1.19	11.1	17.6	1.23	0.0388	<b>2.62</b>	0.13	0.90
<b>Family Characidae</b>										
<i>Astyanax bimaculatus</i>	2587	8.0	1.32	5.4	13.6	1.26	0.0388	<b>2.88</b>	0.07	0.90
<i>Astyanax fasciatus</i>	224	9.1	1.39	7.0	13.2	1.25	0.0307	<b>2.88</b>	0.05	0.94
<i>Acestrorhynchus lacustris</i>	1617	16.5	3.22	10.4	26.4	1.19	0.0062	<b>3.26</b>	0.01	0.96
<i>Galeocharax knerii</i>	1027	14.8	3.25	7.4	24.0	1.20	0.0091	<b>3.18</b>	0.07	0.97
<i>Salminus maxillosus</i>	503	37.6	12.59	12.6	77.3	1.20	0.0110	<b>3.16</b>	0.01	0.99
<i>Brycon orbignyanus</i>	102	26.3	11.50	11.3	62.5	1.21	0.0129	<b>3.14</b>	0.02	0.99
<i>Serrasalmus marginatus</i>	2468	13.1	3.19	4.4	24.1	1.22	0.0162	<b>3.23</b>	0.01	0.98
<i>Pygocentrus nattereri</i>	41	29.4	2.23	24.0	33.3	1.17	0.0269	3.23	0.16	0.91
<i>Serrasalmus spilopleura</i>	122	11.6	4.15	5.1	25.0	1.22	0.0149	<b>3.34</b>	0.03	0.99
<i>Myloplus levis</i>	79	15.4	5.55	5.4	30.1	0.87	0.0517	2.95	0.05	0.98
<b>Family Curimatidae</b>										
<i>Cyphocharax modesta</i>	368	11.0	2.16	6.5	15.5	1.26	0.0353	<b>2.90</b>	0.03	0.96
<i>Cyphocharax nagelii</i>	94	10.3	2.05	6.7	14.4	1.26	0.0346	<b>2.86</b>	0.05	0.97
<i>Leporellus vittatus</i>	288	12.4	2.36	9.0	24.5	1.23	0.0243	2.92	0.05	0.92
<i>Leporinus elongatus</i>	370	23.4	6.93	8.6	51.7	1.22	0.0219	3.00	0.03	0.97
<i>Leporinus friderici</i>	954	18.7	4.61	7.9	37.0	1.24	0.0152	<b>3.14</b>	0.01	0.98
<i>Leporinus obtusidens</i>	318	21.8	6.38	7.5	40.0	1.23	0.0171	<b>3.12</b>	0.02	0.98
<i>Leporinus octofasciatus</i>	166	13.6	3.35	9.1	23.5	1.27	0.0152	<b>3.13</b>	0.03	0.98
<i>Leporinus amblyrhynchus</i>	70	10.9	0.84	9.2	13.6	1.24	0.0747	<b>2.40</b>	0.10	0.89
<i>Leporinus lacustris</i>	57	12.7	2.87	7.0	16.8	1.24	0.0201	<b>3.19</b>	0.06	0.98
<i>Prochilodus lineatus</i>	3719	29.1	6.59	5.9	54.2	1.23	0.0212	<b>3.07</b>	0.01	0.97
<i>Schizodon borellii</i>	1066	21.3	3.42	8.2	38.4	1.19	0.0182	<b>3.08</b>	0.02	0.94
<i>Schizodon nasutus</i>	301	19.6	5.13	10.2	33.6	1.22	0.0155	<b>3.04</b>	0.02	0.99
<i>Schizodon altoparanae</i>	330	20.0	3.04	8.9	31.2	1.20	0.0101	<b>3.26</b>	0.05	0.92
<i>Schizodon cf. platae</i>	52	29.8	2.69	25.4	35.7	1.19	0.0368	2.78	0.16	0.86
<i>Steindachnerina insculpta</i>	5833	9.6	1.14	6.2	14.4	1.26	0.0268	2.99	0.02	0.86
<b>Family Hemiodontidae</b>										
<i>Apareiodon affinis</i>	164	11.2	1.05	7.4	13.5	1.19	0.0720	<b>2.42</b>	0.09	0.81
<i>Hemiodus orthonops</i>	38	20.9	1.87	16.7	24.6	1.23	0.0238	2.93	0.22	0.83
<i>Parodon tortuosus</i>	65	11.9	1.68	9.2	15.5	1.21	0.0362	<b>2.81</b>	0.10	0.93
<b>Family Erythrinidae</b>										
<i>Hoplias aff. malabaricus</i>	604	25.8	6.49	9.6	48.6	1.22	0.0136	<b>3.12</b>	0.03	0.96
<b>Family Cynodontidae</b>										
<i>Rhaphiodon vulpinus</i>	1420	36.7	6.70	13.9	64.0	1.12	0.0031	<b>3.23</b>	0.02	0.95
<b>Family Gymnotidae</b>										
<i>Gymnotus carapo*</i>	70	23.7	4.33	18.5	42.0		0.0031	3.02	0.10	0.93
<b>Family Sternopygidae</b>										
<i>Eigenmannia trilineata*</i>	36	22.6	4.00	16.5	31.5		0.0055	<b>2.66</b>	0.15	0.90
<i>Eigenmannia virescens*</i>	124	22.7	4.26	12.7	37.5		0.0172	<b>2.45</b>	0.07	0.90
<b>Family Apterontidae</b>										
<i>Apterontus albifrons</i>	26	28.4	7.20	20.3	47.5	1.65	0.0029	3.07	0.22	0.89
<i>Apterontus sp.</i>	190	25.0	3.39	16.7	31.7	1.08	0.0088	2.65	0.08	0.87
<b>Family Rhamphichthyidae</b>										
<i>Rhamphichthys rostratus*</i>	37	72.8	16.59	30.0	103.0		0.0020	2.94	0.16	0.90
<b>Family Doradidae</b>										
<i>Pterodoras granulosus</i>	1651	25.7	9.55	5.1	53.2	1.26	0.0365	<b>2.88</b>	0.01	0.97
<i>Rhinodoras dorbignyi</i>	164	12.1	2.65	7.1	19.5	1.22	0.0203	3.06	0.06	0.93
<i>Trachydoras paraguayensis</i>	2491	8.3	0.95	5.3	12.0	1.25	0.0339	<b>2.48</b>	0.21	0.89
<b>Family Auchenipteridae</b>										
<i>Auchenipterus nuchalis</i>	12867	16.6	2.54	8.4	27.0	1.20	0.0150	<b>2.96</b>	0.01	0.89
<i>Parauchenipterus galeatus</i>	3578	13.4	1.92	6.6	22.0	1.22	0.0337	2.99	0.02	0.90
<b>Family Ageneiosidae</b>										
<i>Ageneiosus brevifilis</i>	32	32.4	9.72	17.5	53.2	1.16	0.0200	2.98	0.07	0.98
<i>Ageneiosus ucayalensis</i>	2476	17.7	2.91	11.1	32.4	1.22	0.0107	<b>2.97</b>	0.02	0.93
<i>Ageneiosus valenciennesi</i>	88	22.3	4.75	10.4	31.6	1.21	0.0067	<b>3.19</b>	0.06	0.97

continued

Table 1. (continued)

Family/Species	Standard Length					Length-weight relationship				
	n	mean	s.d.	min	max	TL/LS	a	b	s.e.(b)	r <sup>2</sup>
<b>Family Pimelodidae</b>										
<i>Iheringichthys labrosus</i>	3215	15.3	3.22	7.3	26.2	1.27	0.0087	3.23	0.0	0.97
<i>Pimelodella</i> spp.	128	11.7	1.53	7.4	16.0	1.26	0.0275	2.68	0.08	0.91
<i>Pimelodus maculatus</i>	1474	19.7	5.51	8.3	36.0	1.25	0.0161	3.08	0.01	0.98
<i>Pmelodus ornatus</i>	31	23.5	5.75	9.0	38.5	1.19	0.0119	3.145	0.09	0.98
<i>Pimelodus</i> spp.	119	14.9	4.93	8.0	28.6	1.24	0.0110	3.18	0.07	0.94
<i>Zungaro zungaro</i>	54	26.8	9.50	16.0	52.0	1.21	0.0160	3.16	0.07	0.98
<i>Rhamdia</i> spp.	33	16.8	4.20	10.2	25.2	1.22	0.0234	2.92	0.13	0.94
<i>Hemisorubim platyrhynchos</i>	184	30.8	6.51	17.9	52.5	1.19	0.0131	3.02	0.04	0.96
<i>Paulicea luetkeni</i>	350	43.1	11.51	18.0	97.5	1.21	0.0281	2.94	0.03	0.97
<i>Pseudoplatystoma corruscans</i>	347	59.4	15.00	22.3	114.0	1.15	0.0059	3.16	0.03	0.97
<i>Sorubim lima</i>	79	34.8	7.27	16.8	50.4	1.16	0.0109	2.94	0.08	0.95
<i>Pinirampus pirinampu</i>	466	38.6	11.62	14.9	77.4	1.24	0.0072	3.18	0.02	0.98
<b>Family Hypophthamidae</b>										
<i>Hypophthalmus edentatus</i>	10154	26.6	3.79	9.6	44.6	1.19	0.0075	3.08	0.01	0.95
<b>Family Loricariidae</b>										
<i>Hypostomus</i> spp.	1403	14.7	5.33	5.8	43.6	1.35	0.0290	2.99	0.01	0.97
<i>Megalancistrus aculeatus</i>	76	27.9	8.29	8.6	44.6	1.33	0.0239	3.10	0.05	0.98
<i>Rhinelepis aspera</i>	173	30.5	3.18	20.0	37.2	1.28	0.0873	2.70	0.07	0.89
<i>Loricaria proluxa</i>	28	27.5	9.75	11.0	45.2	1.16	0.0032	3.23	0.06	0.99
<i>Loricaria</i> sp.	1241	21.3	2.84	10.0	27.2	1.14	0.0020	3.35	0.02	0.96
<i>Loricariichthys</i> sp.	4905	20.9	2.66	11.8	33.0	1.14	0.0030	3.25	0.01	0.93
<i>Loricariichthys platymetopon</i>	225	19.4	3.90	11.5	29.0	1.16	0.0048	3.17	0.04	0.96
<b>Family Sciaenidae</b>										
<i>Pachyurus bonariensis</i>	25	16.8	2.48	12.0	22.6	1.23	0.0304	2.82	0.19	0.90
<i>Plagioscion squamosissimus</i>	7200	22.6	4.40	6.8	47.9	1.22	0.0157	3.09	0.01	0.97
<b>Family Cichlidae</b>										
<i>Crenicichla nierderleinii</i>	343	12.7	1.84	7.6	18.5	1.19	0.0330	2.74	0.04	0.92
<i>Crenicichla haroldoi</i>	43	10.9	1.67	8.9	16.1	1.21	0.0370	2.71	0.12	0.93
<i>Crenicichla britskii</i>	64	10.8	1.87	8.1	16.1	1.21	0.1039	2.34	0.12	0.88
<i>Satanoperca pappaterra</i>	153	12.2	2.76	5.9	14.4	1.27	0.0284	3.13	0.03	0.98
<b>Family Archiridae</b>										
<i>Catathyridium jenynsii</i>	57	14.5	4.88	5.5	25.0	1.24	0.0354	3.09	0.06	0.98

( $p < 0.001$ ). The  $b$  values ranged from 2.34 for *Crenicichla britskii* to 3.35 for *Loricaria* sp.

Fig. 2 shows the distribution of exponents of the weight-length relationship for the 72 species under analysis. A tendency towards normality for this type of distribution may be noted, as has been affirmed by Carlander (1969). However, asymmetry to the left is determined ( $G_1 = -0.9877$ ) in the mesocurtic curve ( $G_2 = 0.5652$ ). The average value of  $b$  was 2.986 (s.d.=0.230) and not significantly different from 3 (t-test,  $df=71$ ,  $p=0.05$ ). This value corresponds to that found by Petrakis and Stergiou (1995) and Entsua-Mensah et al. (1995) for average




Fig. 2. Distribution of  $b$  values of the length-weight relationship.

values of  $b$  they analyzed (2.987 and 2.98, respectively). On the other hand, Torres (1992) recorded an average value of  $b = 3.112$  for 40 species of Lake Kariba, Africa, which is significantly higher than 3.

In the analysis of each species it was established that the majority (52) had values of  $b$  significantly different from 3. This proves the indispensable use of the allometric condition factor in these cases. According to Braga (1986), in the case of these species,  $k$  does not vary with fish length. This is a basic premise emphasized by Weatherley and Gill (1987). The correct interpretation of the parameters resulting from the length-weight relationship of the species will disclose information that is useful to the study of fishing biology and management. In this case it should be emphasized that the application of the results of the above-mentioned regressions should be restricted to specimens of the respective species that present standard length within the range of values described in this study.

### Acknowledgments

Thanks are due to the Nucleus for Research in Limnology, Ichthyology and Aquaculture (NUPELIA) for logistical support; to Itaipu Binacional for the financial support; to Jaime Lopes Pereira for drawing the graphs; to librarian

Maria Salette Ribelatto Arita for bibliographical assistance; and to Thomas Bonnici and Lucasz Glowacki for the translation.

### References

- Almeida, F.P., D.L. Lhartley and J. Burnett. 1995. Length-weight relationships and sexual maturity of goosfish of the Northeast coast of the United States. *N. Am. J. Fish. Manage.* 15:14-25.
- Benedito-Cecilio, E. 1994. Dominância, uso do ambiente e associações interespecíficas na ictiofauna do reservatório de Itaipu e alterações decorrentes do represamento. São Carlos, UFSCar, 1994, 173 p. Tese (doutorado em ecologia e recursos naturais). UFSCar.
- Braga, F.M. de S. 1986. Estudo entre fator de condição e relação peso/comprimento para alguns peixes marinhos. *Rev. Brasil. Biol.* 46 (2): 339-346.
- Carlander, K.D. 1969. Handbook of freshwater fishery biology. Iowa State University Press, Ames.
- Entsua-Mensah, M., A. Osci-Abunyewa and M.L.D. Palomares. 1995. Length-weight relationships of fishes from tributaries of the Volta river, Ghana. Part 1. Analysis of pooled data sets. *Naga. ICLARM Q.* 18(1/2): 36-38.
- Gulland, J. A. 1983. Fish stock assessment: a manual of basic methods. v. 1 (series on Food and Agriculture). FAO/Wiley and Sons, New York. 223 p.
- Kulbicki, M., G. Mou Tham, P. Thollot and L. Wantiez. 1993. Length-weight relationship of fish from the Lagoon of New Caledonia. *Naga, ICLARM Q.* 16(2/3): 26-30.
- Pauly, D. 1993. FishByte Editorial. *Naga, ICLARM Q.* 16(2/3):26.
- Petrakis, G. and K.I. Stergiou. 1995. Weight-length relationships for 33 fish species in Greek waters. *Fish. Res.* 21: 465-469.
- Stergiou, K.I. and C.Y. Politou. 1995. Biological parameters, body length-weight and length-height relationships for various species in Greek waters. *Naga, ICLARM Q.* 18(2):42-45.
- Torres Jr., F.S.B. 1992. Length-weight relationships of Lake Kariba fishes. *Naga, ICLARM Q.* 15(4):42-43.
- Vazzoler, A.E.A. de M. 1996. Biologia da reprodução de peixes teleosteos: teoria e prática. Maringá: EDUEM. São Paulo: SBI. 169 p.
- Weatherley A.H. and H.S. Gill. 1987. The biology of fish growth. London: Academic Press, 443 p.
- E. BENEDITO-CECILIO and A.A. AGOSTINHO** are from the *Research Nucleus in Limnology, Ichthyology and Aquaculture, Department of Biology of the Universidade Estadual de Maringá (Paraná, Brazil)* while **R.C. CARNELÓS-MACHADO VELHO** is from the *Biological Sciences Course of the Universidade Estadual de Maringá (Paraná, Brazil)*.

